



TITLE:

# Relationship between depression and risk of malnutrition among community-dwelling young-old and old-old elderly people.

AUTHOR(S):

Yoshimura, Kazuya; Yamada, Minoru; Kajiwar, Yuu; Nishiguchi, Shu; Aoyama, Tomoki

---

CITATION:

Yoshimura, Kazuya ...[et al]. Relationship between depression and risk of malnutrition among community-dwelling young-old and old-old elderly people.. Aging & mental health 2012, 17(4): 456-460

ISSUE DATE:

2012-11

URL:

<http://hdl.handle.net/2433/189833>

RIGHT:

This is an Accepted Manuscript of an article published by Taylor & Francis Group in "Aging and Mental Health" on 11/26/2012, available online at: <http://www.tandfonline.com/10.1080/13607863.2012.743961>; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

**Relationship between depression and risk of malnutrition among  
community-dwelling young-old and old-old elderly people**

**Authors**

Kazuya Yoshimura, Department of Physical Therapy, Human Health Sciences,  
Graduate School of Medicine, Kyoto University, Kyoto, Japan.

Minoru Yamada, Department of Physical Therapy, Human Health Sciences,  
Graduate School of Medicine, Kyoto University, Kyoto, Japan.

Yuu Kajiwarra, Department of Physical Therapy, Human Health Sciences,  
Graduate School of Medicine, Kyoto University, Kyoto, Japan.

Shu Nishiguchi, Department of Physical Therapy, Human Health Sciences,  
Graduate School of Medicine, Kyoto University, Kyoto, Japan.

Tomoki Aoyama, Department of Physical Therapy, Human Health Sciences,  
Graduate School of Medicine, Kyoto University, Kyoto, Japan.

21

22 Corresponding author:

23 Kazuya Yoshimura

24 Department of Physical Therapy, Human Health Sciences, Graduate School of

25 Medicine, Kyoto University, Kyoto, Japan.

26 53 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

27 E-mail: [yoshimura.kazuya.76w@st.kyoto-u.ac.jp](mailto:yoshimura.kazuya.76w@st.kyoto-u.ac.jp)

28 Phone: +81-75-751-3935

29 Fax: +81-75-751-3909

30

31

32 Title

33 Relationship between depression and risk of malnutrition among  
34 community-dwelling young-old and old-old elderly people

35

36 Journal

37 Aging & Mental Health

38

39 Abstract

40 Objectives: The present study explores the association between nutritional  
41 status and depression among healthy community-dwelling young-old (aged 65–  
42 74) and old-old elderly (aged 75 and older).

43

44 Method: A cross-sectional design was implemented. A total of 274  
45 community-dwelling older individuals (142 young-old; 132 old-old) were  
46 assessed using the Geriatric Depression Scale (GDS), Short-Form  
47 Mini-Nutritional Assessment (MNA-SF), and Life-Space Assessment (LSA).  
48 Logistic regression analysis was used to determine if depression was  
49 independently associated with risk of malnutrition, stratified by age (young-old vs.  
50 old-old).

51

Results: In the logistic regression model for young-old, being at risk of malnutrition ( $\text{MNA-SF} \leq 11$ ) was strongly associated with depression ( $\text{GDS} \geq 5$ ) (likelihood ratio = 6.26; 95% confidence interval [CI]: 1.91–20.49). In contrast, in the old-old group, the model was not statistically significant.

Conclusion: Depression and nutritional status were strongly correlated in young-old but not in old-old community-dwelling elderly. This study reveals that not only the factors correlated with but also the symptoms of depression may vary among different age stratifications of the elderly.

Keywords: Depression, Nutritional Status, Young-old

## **Relationship between depression and risk of malnutrition among community-dwelling young-old and old-old elderly people**

### **Introduction**

In an aging population, depression is a serious public health issue due to its societal burden and association with various factors. The total cost of depression in Japan in 2005 was estimated to be 2.0 trillion yen (USD 1 = JPY 78: October 2012) (Sado et al., 2011), which includes all direct, morbidity, and mortality costs. Further, studies in a variety of settings have shown that depression is strongly associated with low function and poor quality of life, increase in the use of health services, late-life suicide tendency, and excess mortality (Callahan, Hui, Nienaber, Musick, & Tierney, 1994; Cuijpers & Smit, 2002; Geerlings, Beekman, Deeg, Twisk, & Van Tilburg, 2002; Koenig, Shelp, Goli, Cohen, & Blazer, 1989; Nyunt, Lim, Yap, & Ng, 2012; Turvey et al., 2002; Van der Weele, Gussekloo, De Waal, De Craen, & Van der Mast, 2009; Wada et al., 2005). A systematic review of 34 community-based studies reported that the prevalence of depression in the elderly is as high as 35% (Beekman, Copeland, & Prince, 1999; Woo et al., 1994). In terms of the type of depression, major depression is relatively rare and minor depression more common (Beekman, Copeland, & Prince, 1999). In

84 addition, clinicians often fail to diagnose and treat depression in the elderly, and  
85 elderly individuals are reluctant to report depressive symptoms (Georgotas,  
86 Cooper, Kim, & Hapworth, 1983; Lyness et al., 1995; Webber et al., 2005).  
87 Therefore, depression in the elderly has often been unrecognized, and it is  
88 important to clarify the components of depression to improve the sensitivity of  
89 screening methods.

90  
91 Many variables, including gender, marital status, cognitive status, activities of  
92 daily living and independent activities of daily living limitations, and social  
93 engagement, are associated with increased depression (Anstey, Von Sanden,  
94 Sargent-Cox, & Luszcz, 2007; Glass, Mendes De Leon, Bassuk, & Berkman,  
95 2006). Meanwhile, weight loss and loss of appetite are the main factors that  
96 mediate the relationship between depression and nutritional status (Rubenstein,  
97 Harker, Salvà, Guigoz, & Vellas, 2001). Recently, multiple studies have indicated  
98 an association between depression and nutritional status in various settings  
99 such as outpatient clinics and institutions and in the community (Cabrera, Mesas,  
100 Garcia, & Andrade, 2007; Kaburagi et al., 2011; Smoliner et al., 2009; Wilson,  
101 Vaswani, Liu, Morley, & Miller, 1998). In these settings, depression is an  
102 independent predictor of nutritional health and a major cause of weight loss  
103 (Chen, Chang, Chyun, & McCorkle, 2005; Morley & Kraenzle, 1994; Thompson

& Morris, 1991). Conversely, better diet quality is beneficial for preventing and improving depressive symptoms (Akbaraly et al., 2009; Lin & Su, 2007). The relationship between depression and nutritional status is interactive and further investigation is still needed. In this regard, the aging society needs to be considered. The word 'elderly' is very broad, encompassing all individuals aged older than 60 or 65 years. Age may alter the relationship among the various variables associated with depression. For example, the young-old (age 65–74 years) and old-old (age 75 years and above) have different predispositions regarding various aspects of both depression and nutritional status (Kaburagi et al., 2011; Kondo, Kazama, Suzuki, & Yamagata, 2008). Older elderly are more likely to experience frailty, physical illness, bereavement, and loneliness, which are risk factors contributing to depression (Bruce, 2002; Blazer, 2002). Investigating age-group differences may help in clarifying the correlation between depression and nutritional status and in developing precise, age-stratified interventions.

The purpose of the present study was to analyse the association between depression and nutritional status in different age groups of non-functionally-impaired, community-dwelling elderly.



124

## 125    Methods

## 126    [Participants]

127    Study participants were recruited through advertisements in the local press and  
128    at local healthcare events from April 2011 to June 2012. A total of 274  
129    community-dwelling older individuals volunteered in the study. The inclusion  
130    criteria were an age of 65 years or older, living in the community, and being able  
131    to walk independently with or without a cane. The exclusion criteria ensured that  
132    none of the participants had any indications of the following health problems: (a)  
133    symptomatic cardiovascular disease, (b) neurological and orthopaedic disorders,  
134    (c) peripheral neuropathy of the lower extremities, and (d) severe arthritis.

135

136    This study was approved by the Ethical Review Board of Kyoto University  
137    Graduate School of Medicine, Kyoto, Japan.

138

## 139    [Variables]

140    The following variables were collected using a questionnaire.

141

142    Depression was screened for by the 15-item Geriatric Depression Scale (GDS)  
143    (Yesavage, 1988), a validated and reliable self-report scale that detects

depression in elderly people. Scores range from 0 to 15. We used a cut-off of 4/5, which is a recommended indicator of depression in Japanese populations (Murata, Kondo, Hirai, Ichida, & Ojima, 2008; Yamazaki, Nakano, Saito, & Yasumura, 2012).

Nutritional status was assessed with the Short-Form Mini-Nutritional Assessment (MNA-SF) (Rubenstein et al., 2001). MNA-SF includes 6 items dealing with loss of appetite, weight loss, mobility, stress or illness, dementia or depression, and body mass index (BMI). Scores range from 0 to 14. A score of 12 or above indicates satisfactory nutritional status, a score of 8 to 11 implies risk of malnutrition, and a score less than 7 suggest malnourishment.

Life-space mobility was assessed by the Life-Space Assessment (LSA) (Baker, Bodner, & Allman, 2003), a questionnaire that measures the spatial extent of individuals in a given month. The LSA takes into account the frequency of travel to different life-space levels (bedroom, driveway, within neighbourhood, outside neighbourhood but within town, and out of town), and whether personal or technical assistance was required to get to those levels. The composite scores range from 0 to 120.

## [Statistical Analysis]

Statistical analysis was carried out with the software package SPSS 20.0 (SPSS Inc., Chicago, IL). Relationships between MNA-SF items and depression were evaluated with chi-square tests. Multivariate analysis was performed to examine the association between depression and risk of malnutrition. Step-up logistic regression analysis was used to determine if depression ( $GDS \geq 5$ ) was independently associated with risk of malnutrition ( $MNA-SF \leq 11$ ). In Model I, all of the participants were analysed. Model II contained only young-old participants, and Model III contained only old-old participants. Demographic factors (age, gender, and BMI) and LSA were adjusted in each model. An acceptable level of statistical significance was considered to be a p value of  $< .05$ .

## Results

Participant characteristics are shown in Table 1. The mean age was 74.33 (SD 4.72) years, and 185 participants (67.5%) were female. Classifying participants by age, 142 were young-old (51.8%) and 132 participants were old-old (48.2%). Fifty-nine participants (21.5%) were depressed. Seventy-seven (28.1%) were at risk of malnutrition (including 1 participant determined to be malnourished), and the others were well nourished. Old-old participants had a higher risk of

malnutrition than young-old ones, but no significant differences were found for depression. The nutritional characteristics of the participants are shown in Table 2. Among the young-old, there was a trend that depressed participants were more likely to have a loss of appetite within the past 3 months than the non-depressed participants, but this difference was not statistically significant ( $p = .075$ ). There was no trend toward significance in the old-old group ( $p = .502$ ).

In the logistic regression model for young-old (Model II), being at risk of malnutrition ( $\text{MNA-SF} \leq 11$ ) was strongly associated with depression ( $\text{GDS} \geq 5$ ) (likelihood ratio = 6.74; 95% confidence interval [CI]: 2.11–21.51) independent from the control variables, while this association was not found in Models I and III (Table 3).

## Discussion

The present study found a correlation between GDS and MNA-SF for young-old individuals, but not for old-old individuals or for the two groups combined. Multiple studies conclude that depression and malnutrition are related, but that the influence of age on the variables differs. Previous studies reported that depression is an independent predictor of malnutrition or nutritional risk even

after adjusting for social and educational factors in young-old elderly adults but not old-old elderly adults (Cabrera et al., 2007; Callen and Wells, 2005). These differences could arise because depressive symptoms in the elderly have different clinical features along the age spectrum from young-old to old-old (Mehta et al., 2008). For example, old-old elderly may suffer from a higher prevalence of disability or medical illnesses (Chou & Chi, 2005). Having a chronic disease is a variable that independently influences depression (Schoevers et al., 2000). The relationship between depression and malnutrition needs to be further examined, with physical, mental, and social status taken into consideration.

Various studies have reported a positive relationship between depression and nutritional status, and they often associate depression with loss of appetite or weight loss (Akbaraly et al., 2009; Davison & Kaplan, 2012). However, the rates of depressed participants in the present study with loss of appetite or weight loss were 13.6% and 30.5% in the young-old and old-old, respectively, and were relatively low. Only the relationship between depression and loss of appetite in young-old elderly showed a trend toward significance, and other relationships (between depression and weight loss in young-old and between depression and loss of appetite or weight loss in old-old) did not. Callen and Wells (2005)

reported that in old-old elderly, depression is not a predictor of weight loss or low BMI when adjusting for social, physical, and economic factors, a finding which is in agreement with our results. The relationship between depression and loss of appetite may have been the principal reason behind the result obtained in the present multivariate analysis, that is, the positive relationship between depression and malnutrition.

Our study has several limitations. The cross-sectional design prevents us from making causal inferences. We also did not assess socioeconomic and educational status and social support; the possibility of these being confounding factors cannot be denied. Despite these limitations, this study reveals that the factors correlated with depression could vary among different age groups of elderly and suggests that depression and nutritional status are correlated more strongly in young-old than old-old elderly. Future studies should focus on clarifying the causal relationship, consider the age of subjects and assess nutritional status, social status, etc., for better understanding of depression.

## Reference

Akbaraly, T.N., Brunner, E.J., Ferrie, J.E., Marmot, M.G., Kivimaki, M., & Singh-Manoux, A. (2009). Dietary pattern and depressive symptoms in middle

- 244 age. *British Journal of Psychiatry*, 195, 408-413
- 245
- 246 Anstey, K.J., Von Sanden, C., Sargent-Cox, K., & Luszcz, M.A. (2007).
- 247 Prevalence and risk factors for depression in a longitudinal, population-based
- 248 study including individuals in the community and residential care. *American*
- 249 *Journal of Geriatric Psychiatry*, 15, 497-505
- 250
- 251 Baker, P.S., Bodner, E.V., & Allman, R.M. (2003). Measuring Life-Space Mobility
- 252 in Community-Dwelling Older Adults. *Journal of the American Geriatrics Society*,
- 253 51, 1610-1614
- 254
- 255 Beekman, A.T.F., Copeland, J.R.M., & Prince, M.J. (1999). Review of community
- 256 prevalence of depression in later life. *British Journal of Psychiatry*, 174, 307-311
- 257
- 258 Bruce, M.L. (2002). Psychosocial risk factors for depressive disorders in late life.
- 259 *Biological Psychiatry*, 52, 175-184
- 260
- 261 Blazer, DG. (2002). *Depression in Later Life*. New York: Springer.
- 262
- 263 Cabrera, M.A.S., Mesas, A.E., Garcia, A.R.L., & de Andrade, S.M. (2007).

- 264     Malnutrition and Depression among Community-dwelling Elderly People.  
265     *Journal of the American Medical Directors Association*, 8, 582-584  
266
- 267     Callahan, C.M., Hui, S.L., Nienaber, N.A., Musick, B.S., & Tierney, W.M. (1994).  
268     Longitudinal study of depression and health services use among elderly primary  
269     care patients. *Journal of the American Geriatrics Society*, 48, 833-838  
270
- 271     Callen, B.L. & Wells, T.J. (2005). Screening for nutritional risk in  
272     community-dwelling old-old. *Public Health Nursing*, 22, 138-146  
273
- 274     Chen, C.C.-H., Chang, C.-K., Chyun, D.A., & McCorkle, R. (2005). Dynamics of  
275     nutritional health in a community sample of American elders: A multidimensional  
276     approach using Roy Adaptation Model. *Advances in Nursing Science*, 28,  
277     376-389  
278
- 279     Chou, K.-L. & Chi, I. (2007). Prevalence and correlates of depression in Chinese  
280     oldest-old. *International Journal of Geriatric Psychiatry*, 20, 41-50  
281
- 282     Cuijpers, P. & Smit, F. (2002). Excess mortality in depression: A meta-analysis of  
283     community studies. *Journal of Affective Disorders*, 72, 227-236.



284

285 Davison, K.M. & Kaplan, B.J. (2012). Food intake and blood cholesterol levels of  
286 community-based adults with mood disorders. *BMC Psychiatry*. Retrieved from  
287 [http://dx.doi.org/ 10.1186/1471-244X-12-10](http://dx.doi.org/10.1186/1471-244X-12-10)

288

289 Geerlings, S.W., Beekman, A.T., Deeg, D.J., Twisk, J.W., & Van Tilburg, W.  
290 (2002). Duration and severity of depression predict mortality in older adults in the  
291 community. *Psychological Medicine*, 32, 609-618.

292

293 Georgotas, A., Cooper, T., Kim, M., & Hapworth, W. (1983). The treatment of  
294 affective disorders in the elderly. *Psychopharmacology Bulletin*, 19, 226-237

295

296 Glass, T.A., Mendes De Leon, C.F., Bassuk, S.S., & Berkman, L.F. (2006). Social  
297 engagement and depressive symptoms in late life: Longitudinal findings. *Journal*  
298 *of Aging and Health*, 18, 604-628

299

300 Kaburagi, T., Hirasawa, R., Yoshino, H., Odaka, Y., Satomi, M., Nakano, M., ...,  
301 Sato, K. (2011). Nutritional status is strongly correlated with grip strength and  
302 depression in community-living elderly Japanese. *Public Health Nutrition*, 14,  
303 1893-1899

304

305 Koenig, H.G., Shelp, F., Goli, V., Cohen, H.J., & Blazer, D.G. (1989). Survival  
306 and health care utilization in elderly medical inpatients with major depression.  
307 *Journal of the American Geriatrics Society*, 37, 599-606.

308

309 Kondo, N., Kazama, M., Suzuki, K., & Yamagata, Z. (2008). Impact of mental  
310 health on daily living activities of Japanese elderly. *Preventive Medicine*, 46,  
311 457-462

312

313 Lin, P.-Y. & Su, K.-P. (2007). A meta-analytic review of double-blind,  
314 placebo-controlled trials of antidepressant efficacy of omega-3 fatty acids.  
315 *Journal of Clinical Psychiatry*, 68, 1056-1061

316

317 Lyness, J.M., Cox, C., Curry, J., Conwell, Y., King, D.A., & Caine, E.D. (1995).  
318 Older age and the underreporting of depressive symptoms. *Journal of the*  
319 *American Geriatrics Society*, 43, 216-221

320

321 Mehta, M., Whyte, E., Lenze, E., Hardy, S., Roumani, Y., Subashan, ...,  
322 Studenski, S. (2008). Depressive symptoms in late life: Associations with apathy,  
323 resilience and disability vary between young-old and old-old. *International*

324 *Journal of Geriatric Psychiatry*, 23, 238-243

325

326 Morley, J.E. & Kraenzle, D. (1994). Causes of weight loss in a community  
327 nursing home. *Journal of the American Geriatrics Society*, 42, 583-585

328

329 Murata, C., Kondo, K., Hirai, H., Ichida, Y., & Ojima, T. (2008). Association  
330 between depression and socio-economic status among community-dwelling  
331 elderly in Japan: The Aichi Gerontological Evaluation Study (AGES). *Health and*  
332 *Place*, 14, 406-414

333

334 Nyunt, M.S., Lim, M.L., Yap, K.B., & Ng, T.P. (2012). Changes in depressive  
335 symptoms and functional disability among community-dwelling depressive older  
336 adults. *International Psychogeriatrics* (Epub ahead of print). Retrieved from  
337 <http://dx.doi.org/10.1017/S1041610212000890>

338

339 Rubenstein, L.Z., Harker, J.O., Salvà, A., Guigoz, Y., & Vellas, B. (2001).  
340 Screening for undernutrition in geriatric practice: Developing the Short-Form  
341 Mini-Nutritional Assessment (MNA-SF). *Journals of Gerontology - Series A*  
342 *Biological Sciences and Medical Sciences*, 56, M366-M372

343

344 Sado, M., Yamauchi, K., Kawakami, N., Ono, Y., Furukawa, T.A., Tsuchiya, M.,  
345 ..., Kashima, H. (2011). Cost of depression among adults in Japan in 2005.  
346 *Psychiatry and Clinical Neurosciences*, 65, 442-450.  
347  
348 Schoevers, R.A., Beekman, A.T.F., Deeg, D.J.H., Geerlings, M.I., Jonker, C., &  
349 Van Tilburg, W. (2000). Risk factors for depression in later life; results of a  
350 prospective community based study (AMSTEL). *Journal of Affective Disorders*,  
351 59, 127-137  
352  
353 Smoliner, C., Norman, K., Wagner, K.-H., Hartig, W., Lochs, H., & Pirlich, M.  
354 (2009). Malnutrition and depression in the institutionalised elderly. *Current*  
355 *Opinion in Clinical Nutrition and Metabolic Care*, 11, 1-6  
356  
357 Thompson, M.P. & Morris, L.K. (1991). Unexplained weight loss in the  
358 ambulatory elderly. *Journal of the American Geriatrics Society*, 39, 497-500  
359  
360 Turvey, C.L., Conwell, Y., Jones, M.P., Phillips, C., Simonsick, E., Pearson, J.L.,  
361 & Wallace R. (2002). Risk factors for late-life suicide: A prospective,  
362 community-based study. *American Journal of Geriatric Psychiatry*, 10, 398-406.  
363

- 364 Van der Weele, G.M., Gussekloo, J., De Waal, M.W.M., De Craen, A.J.M., & Van  
365 der Mast, R.C. (2009). Co-occurrence of depression and anxiety in elderly  
366 subjects aged 90 years and its relationship with functional status, quality of life  
367 and mortality. *International Journal of Geriatric Psychiatry*, 24, 595-601  
368
- 369 Wada, T., Ishine, M., Sakagami, T., Kita, T., Okumiya, K., Mizuno, K., ...,  
370 Matsubayashi, K. (2005). Depression, activities of daily living, and quality of life  
371 of community-dwelling elderly in three Asian countries: Indonesia, Vietnam, and  
372 Japan. *Archives of Gerontology and Geriatrics*, 41, 271-280.  
373
- 374 Webber, A.P., Martin, J.L., Harker, J.O., Josephson, K.R., Rubenstein, L.Z., &  
375 Alessi, C.A. (2005). Depression in older patients admitted for postacute nursing  
376 home rehabilitation. *Journal of the American Geriatrics Society*, 53, 1017-1022  
377
- 378 Wilson, M.-M.G., Vaswani, S., Liu, D., Morley, J.E., & Miller, D.K. (1998).  
379 Prevalence and causes of undernutrition in medical outpatients. *American*  
380 *Journal of Medicine*, 104, 56-63  
381
- 382 Woo, J., Ho, S.C., Lau, J., Yuen, Y.K., Chiu, H., Lee, H.C., & Chi, I. (1994). The  
383 prevalence of depressive symptoms and predisposing factors in an elderly

384 Chinese population. *Acta Psychiatrica Scandinavica*, 89, 8-13

385

386 Yamazaki, S., Nakano, K., Saito, E., & Yasumura, S. (2012). Prediction of

387 functional disability by depressive state among community-dwelling elderly in

388 Japan: A prospective cohort study. *Geriatrics and Gerontology International*.

389 Article in Press.

390

391 Yesavage, J.A. (1988). Geriatric Depression Scale. *Psychopharmacology*

392 *bulletin*, 24, 709–711.

393

394

Table 1. Participant characteristics according to age

	All participants		Young-old		Old-old		
	(n = 274)		(age 65–75; n = 142)		(age ≥ 75; n = 132)		
	mean	SD	mean	SD	mean	SD	p value
Age (years)	74.33	4.72	70.61	2.33	78.34	3.07	< .001
Female, %	67.5		66.2		68.9		.699
BMI	22.44	3.26	22.65	3.21	22.23	3.31	.287
LSA (range, 0–120)	87.83	21.08	89.91	20.62	85.59	21.41	.107
MNA-SF score (range, 0–14)	12.32	1.56	12.44	1.50	12.20	1.61	.253
At risk of malnutrition, % (MNA-SF score ≤ 11)†	28.1		21.8		34.8		.022
GDS score (range, 0–15)	2.71	2.74	2.55	2.70	2.89	2.78	.228
Depression, % (GDS score ≥ 5)	21.5		19.7		23.5		.466

BMI: body mass index; LSA: Life-Space Assessment; MNA-SF: Short-Form Mini-Nutrition Assessment; GDS: Geriatric

Depression Scale

† Includes one participant classified as malnourished in the young-old group

p values determined by Mann-Whitney U test or chi-square test

395

396

Table 2. Nutritional characteristics, measured by MNA-SF, of young-old and old-old participants with and without depression

	With depression (GDS ≥ 5)				Without depression (GDS < 5)				
	n (%)				n (%)				p value
MNA-SF items (ranking)†	0	1	2	3	0	1	2	3	
Young-old (n = 142)									
Loss of appetite	1 (3.6)	3 (10.7)	24 (85.7)	-	0 (0)	5 (4.4)	109 (95.6)	-	.075
Weight loss	0 (0)	3 (10.7)	6 (21.4)	19 (67.9)	0 (0)	6 (5.3)	13 (11.4)	95(83.3)	.108
Mobility	0 (0)	-	28 (100)	-	0 (0)	-	114 (100)	-	-
Stress or acute illness	6 (21.4)	-	22 (78.6)	-	7 (6.1)	-	107 (93.9)	-	.022*
Neuropsychological status	2 (7.1)	1 (3.6)	25 (89.3)	-	0 (0)	0 (0)	114 (100)	-	.007**
Body mass index	4 (14.3)	6 (21.4)	7 (25.0)	11 (39.3)	11 (9.6)	16 (14.0)	45 (39.5)	42 (36.8)	1.000
Old-old (n = 132)									



Loss of appetite	0 (0)	4 (12.9)	27 (87.1)	-	0 (0)	9 (8.9)	92 (91.1)	-	.5927
Weight loss	0 (0)	3 (9.7)	6 (19.4)	22 (71.0)	3 (3.0)	6 (5.9)	12 (11.9)	80 (79.2)	.3398
Mobility	0 (0)	-	31 (100)	-	0 (0)	-	101 (100)	-	.399
Stress or acute illness	6 (19.4)	-	25 (80.6)	-	8 (7.9)	-	93 (92.1)		.0760
Neuropsychological status	1 (3.2)	1 (3.2)	29 (93.5)	-	0 (0)	2 (2.0)	99 (98.0)	-	.2461
Body mass index	3 (9.7)	8 (25.8)	6 (19.4)	14 (40.6)	17 (16.8)	21 (20.8)	22 (21.8)	41 (40.6)	.681

MNA-SF: Short-Form Mini-Nutrition Assessment; GDS: Geriatric Depression Scale

† Higher scores indicate better function

Chi-square test: Loss of appetite and neuropsychological status, 0–1 vs. 2; weight loss and body mass index, 0–2 vs. 3; stress or acute

illness, 0 vs. 2

\* p < .05, \*\* p < .01

Table 3. Step-up logistic regression model of variables associated with depression (GDS  $\geq 5$ )

	All participants		Young-old		Old-old	
	(Model I)		(Model II)		(Model III)	
	(n = 274)		(age 65–75; n = 142)		(age $\geq 75$ ; n = 132)	
	Likelihood ratio (95% CI)	p value	Likelihood ratio (95% CI)	p value	Likelihood ratio (95% CI)	p value
Risk of malnutrition (MNA-SF $\leq 11$ )	-	NS	6.738** (2.111–21.510)	.001	-	NS
Age (years)	-	NS	-	NS	-	NS
Gender (male 0; female 1)	-	NS	-	NS	-	NS
BMI	-	NS	1.201* (1.033–1.395)	.017	-	NS
LSA	.985* (.971– .999)	.031	-	NS	-	NS

GDS: Geriatric Depression Scale; MNA-SF: Short-Form Mini-Nutrition Assessment; BMI Body Mass Index; LSA: Life-Space Assessment

NS: not selected

---

\*  $p < .05$ , \*\*  $p < .01$

402